



**QUEEN'S
UNIVERSITY
BELFAST**

In patients with shortened dental arches do removable dental prostheses improve masticatory performance?

Moore, C., & McKenna, G. (2016). In patients with shortened dental arches do removable dental prostheses improve masticatory performance? *Evidence-based dentistry*, 17(4). <https://doi.org/10.1038/sj.ebd.6401204>

Published in:
Evidence-based dentistry

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2016 British Dental Association
This work is made available online in accordance with the publisher's policies.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

For many patients replacement of missing teeth may be indicated to restore oral function, aesthetics and improve quality of life. Whilst rates of edentulism have fallen a partially dentate older population has emerged.^{1,2} Unfortunately, there is a limited amount of evidence and little consensus at present about how best to replace missing teeth in these partially dentate patients.³ One option is utilisation of the Shortened Dental Arch (SDA) concept which aims to provide a functional dentition without the need for a removal partial denture (RPD).⁴ Multiple iterations of the SDA exist based on the number and distribution of occluding pairs of posterior teeth.⁵

The aim of this systematic review was to summarise the impact of RPD provision on objective measures of masticatory function in SDA and extreme SDA patients (eSDA). The review question, inclusion criteria and search strategy were clearly defined and a reasonable publication-date restriction was applied. A total of 8 studies were identified and included. However, sample sizes of the included studies were small, ranging from 8-30.

Based on data included from two studies the effect of RPD provision on comminution appears to be inconclusive for SDA patients. One study reported an improvement in comminution, and the other a deterioration, post-RPD provision. A single study showed a 20% improvement in comminution for eSDA patients provided with a RPD. Two further studies demonstrated a longitudinal increase in comminution for eSDA patients provided with RPDs with a greater number of artificial teeth. A similar relationship was found for chewing function and the number of artificial teeth in two separate studies analysing SDA and eSDA. However, increasing the occlusal table through distal-extension saddles, raises the potential for overload of the denture-bearing area and consequently ridge resorption.^{6,7} In addition, patient compliance with such removal prostheses is often very poor.

This systematic review illustrates the paucity of high quality evidence to answer this research question. There was significant clinical and methodological heterogeneity amongst the small number of included studies. Unfortunately an assessment of the 'risk of bias' was not included. Only two studies analysed participants who did not act as their own controls and neither incorporated randomisation at the allocation stage.

The authors accept that whilst the masticatory ability of SDA and eSDA patients may be lower than that of subjects with a complete dentition, the overall prevalence of chewing complaints is low in this population.⁸⁻¹⁰ However, subjective measures of chewing function and quality of life may be useful tools in determining individual patient need for RPDs. In the first instance, patients with eSDAs could be restored to a longer SDA using fixed prosthodontic options.¹¹ This approach has been demonstrated to be more cost-effective than RPDs whilst also achieving positive impacts on quality of life and nutritional status.¹²⁻¹⁵

Practice points:

- A SDA should provide sufficient functionality for the vast majority of patients. Extending a SDA with a RPD is not recommended.

- Small span fixed prosthodontics could be considered as a first-line treatment for eSDA patients with chewing complaints.

References:

1. Slade GD, Akinkugbe AA, Sanders AE. Projections of U.S. Edentulism prevalence following 5 decades of decline. *J Dent Res.* 2014;93:959-65.
2. Murray JJ. Adult dental health surveys: 40 years on. *Br Dent J.* 2011;211:407-8.
3. Abt E, Carr AB, Worthington HV. Interventions for replacing missing teeth: partially absent dentition. *Cochrane Database Syst Rev.* 2012;(2):CD003814.
4. Käyser AF. Shortened dental arches and oral function. *J Oral Rehabil.* 1981;8:457-62.
5. Faggion CM. The shortened dental arch revisited: from evidence to recommendations by the use of the GRADE approach. *J Oral Rehabil.* 2011;38:940-9.
6. Wyatt CC. The effect of prosthodontic treatment on alveolar bone loss: a review of the literature. *J Prosthet Dent.* 1998;80:362-6.
7. D.L. Sarandha. *Textbook of Complete Denture Prosthodontics.* 1st ed. New Delhi: Jaypee Brothers Medical Publishers. 2008.
8. Sarita PT, Witter DJ, Kreulen CM, Van't Hof MA, Creugers NH. Chewing ability of subjects with shortened dental arches. *Community Dent Oral Epidemiol.* 2003;31:328-34.
9. Witter DJ, De Haan AF, Käyser AF, Van Rossum GM. A 6-year follow-up study of oral function in shortened dental arches. Part II: Craniomandibular dysfunction and oral comfort. *J Oral Rehabil.* 1994;21:353-66.
10. Kanno T, Carlsson GE. A review of the shortened dental arch concept focusing on the work by the Käyser/Nijmegen group. *J Oral Rehabil.* 2006;33:850-62.
11. Jepson N, Allen F, Moynihan P, Kelly P, Thomason M. Patient satisfaction following restoration of shortened mandibular dental arches in a randomized controlled trial. *Int J Prosthodont.* 2003;16:409-14.
12. McKenna G, Allen F, Woods N, O'Mahony D, Cronin M, DaMata C, Normand C. Cost-effectiveness of tooth replacement strategies for partially dentate elderly: a randomized controlled clinical trial. *Community Dent Oral Epidemiol.* 2014;42:366-74.
13. Walter MH, Marré B, Vach K, Strub J, Mundt T, Stark H et al. Management of shortened dental arches and periodontal health: 5-year results of a randomised trial. *J Oral Rehabil.* 2014;41:515-22.
14. McKenna G, Allen PF, O'Mahony D, Flynn A, Cronin M, DaMata C et al. Comparison of functionally oriented tooth replacement and removable partial dentures on the nutritional status of partially dentate older patients: a randomized controlled clinical trial. *J Dent* 2014;42:653-9.
15. Wolfart S, Müller F, Gerß J, Heyedcke G, Marré B, Böning K et al. The randomized shortened dental arch study: oral health-related quality of life. *Clin Oral Investig.* 2014;18:525-33.